

Reverse Data-Processing Theorems


Francesco Buscemi (Nagoya University)

Hong Kong Workshop on Quantum Information and Foundations

Dept. of Computer Science, University of Hong Kong


4 May 2016

what is a data-processing theorem?



- ✓ let $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{Y}$ and $\mathcal{W} : \mathcal{Y} \rightarrow \mathcal{Z}$ be two noisy channels
- ✓ then, the joint distributions $(U, Y) \equiv (U, \mathcal{N}(X))$ and $(U, Z) \equiv (U, \mathcal{W}(Y))$ are such that

$$H(U|Y) \leq H(U|Z) \quad \text{for all initial } (U, X)$$



- ✓ let $\mathcal{B} : \mathcal{X} \rightarrow (\mathcal{Y}, \mathcal{Z})$ be a broadcast channel
- ✓ assume that the final distribution $(U, Y, Z) \equiv (U, \mathcal{B}(X))$ is such that

$$H(U|Y) \leq H(U|Z) \quad \text{for all initial } (U, X)$$

- ✓ can we then conclude that there exists a noisy channel $\mathcal{W} : \mathcal{Y} \rightarrow \mathcal{Z}$ such that $(U, Z) = (U, \mathcal{W}(Y))$ for all initial (U, X) ?
- ✓ no (Körner and Marton, 1977)

Given two noisy channels $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{Y}$ and $\mathcal{N}' : \mathcal{X} \rightarrow \mathcal{Z}$, Körner and Marton (1977) introduce the following definitions:



- ✓ \mathcal{N} is **degradable** into \mathcal{N}' if there exists a noisy channel $\mathcal{W} : \mathcal{Y} \rightarrow \mathcal{Z}$ such that

$$\mathcal{N}' = \mathcal{W} \circ \mathcal{N}$$

- ✓ \mathcal{N} is **less noisy** than \mathcal{N}' if

$$H(U|Y) \leq H(U|Z) \quad \text{for all initial } (U, X)$$

- ✓ \mathcal{N} is **more capable** than \mathcal{N}' if

$$H(X|Y) \leq H(X|Z) \quad \text{for all initial } X$$

- ✓ **fact:** degradable $\xRightarrow{\neq}$ less noisy $\xRightarrow{\neq}$ more capable



Reverse Data-Processing Theorem (FB, Prob. Inf. Trans. 2016)

Given two noisy channels $\mathcal{N} : \mathcal{X} \rightarrow \mathcal{Y}$ and $\mathcal{N}' : \mathcal{X} \rightarrow \mathcal{Z}$,

$$H_{\min}(U|Y) \leq H_{\min}(U|Z) \quad \text{for all initial } (U, X),$$


if and only if \mathcal{N} is degradable in \mathcal{N}' . In other words,

$$\text{degradable} \iff \text{less noisy w.r.t. } H_{\min}.$$

- ✓ $H_{\min}(U|Y) = -\log_2 P_{\text{guess}}(U|Y) = -\log_2 \sum_y \max_u p(u, y)$
- ✓ it also holds **approximately**


finding stochastic dependencies

- ✓ suppose that, given a conditional probability $p(y_1, y_2, \dots, y_N | x)$, we want to find stochastic dependencies between these variables



- ✓ **reverse data-processing theorem**: a path exists if and only if H_{\min} never decreases (equivalently, no path exists if and only if H_{\min} strictly decreases at some point for some initial conditions)
- ✓ **stochastic dependencies follow the “flow” of H_{\min}**
- ✓ partial ordering \implies incomparable paths
- ✓ for example, the figure above is equivalent to the following entropic conditions:


$$\begin{cases} H_{\min}(U|Y_1) \leq H_{\min}(U|Y_2) \leq H_{\min}(U|Y_4) \leq H_{\min}(U|Y_5), & \text{for all initial } (U, X) \\ H_{\min}(U|Y_3) \leq H_{\min}(U|Y_5), & \text{for all initial } (U, X) \\ H_{\min}(U|Y_3) \leq \{H_{\min}(U|Y_1), H_{\min}(U|Y_2), H_{\min}(U|Y_4)\}, & \text{for some initial } (U, X) \end{cases}$$



- ✓ divisibility: $\mathcal{N}_i = \mathcal{W}_i \circ \mathcal{W}_{i-1} \circ \dots \circ \mathcal{W}_1$
- ✓ a dynamical map is divisible if and only if the sequence $\{H_{\min}(U|Y_i)\}_{i \geq 1}$ is non-decreasing for all initial (U, X)
- ✓ namely, **divisibility is equivalent to "no H_{\min} backflow"**
- ✓ the same insight holds also in the **quantum** (Buscemi, Datta; PRA 2016) **approximate** (Jencova; ISIT 2016) case

the case of open quantum systems dynamics

S : system, E : environment, $S + E$: conservative




But the given process need not be collisional to be divisible. **Question:** how to characterize system-environment correlations that do not break divisibility?

the “Pechukas debate” (1994 onwards)


- ✓ experimentally, the initial factorization condition is an approximation (and strong-coupling regimes are of interest)
- ✓ what happens to the reduced dynamics in the presence of initial system-environment correlations?
- ✓ Pechukas (PRL, 1994): “Here we show that complete positivity is an artifact of product initial conditions. In general, reduced dynamics need not be CP”
- ✓ Lindblad (J. Phys. A, 1995) and Alicki (PRL, 1995, comment to Pechukas)
- ✓ Rodriguez-Rosario, Modi, Kuah, Sudarshan (2008): null discord \implies divisibility
- ✓ Shabani–Lidar (2009): null discord \iff divisibility (Erratum 2016)
- ✓ Brodutch, Datta, Modi, Rivas, Rodriguez-Rosario (2013): divisibility $\not\implies$ null discord
- ✓ FB (2014): what follows

- ✓ system and environment start in a correlated state: they have to be considered as one composite system
- ✓ reference–system–environment: $\mathcal{H}_U \otimes \mathcal{H}_S \otimes \mathcal{H}_E$
- ✓ the initial condition is given as a tripartite density operator ρ_{USE}



- ✓ the state ρ_{USE} is "passive" if $H(U|S)_\rho \leq H(U|S')_\sigma$ for all isometries V
- ✓ **first fact:** ρ_{USE} passive $\iff \forall V_{SE}, \exists \mathcal{E}_S^V : \sigma_{US'} = (\text{id}_U \otimes \mathcal{E}_S^V)(\rho_{US})$
- ✓ **second fact:** ρ_{USE} passive $\iff I(U; E|S)_\rho = 0 \implies$ "passive" \equiv "Markovian"
- ✓ the above facts constitute another reverse data-processing theorem

- ✓ no reference system; instead, given is a family $\mathcal{S} = \{\rho_{SE} : \rho_{SE} \in \mathcal{S}\}$ of possible initial joint system–environment states



- ✓ in the above diagram, ρ_{SE} is a generic element of \mathcal{S} (i.e., the condition must hold for all $\rho_{SE} \in \mathcal{S}$)
- ✓ if the above diagram holds for all isometries V , we say that the family \mathcal{S} is **Markovian**
- ✓ example: the **initial factorization condition**, i.e., $\mathcal{S} = \{\rho_S \otimes \bar{\xi}_E : \rho_S \text{ any state of } S\}$ for some fixed environment state $\bar{\xi}_E$, defines a Markovian family

- ✓ how to connect the tripartite with the bipartite scenario?
- ✓ steering: $\rho_{SE}^{\Pi} = \frac{\text{Tr}_U[\rho_{USE} (\Pi_U \otimes \mathbf{1}_{SE})]}{\text{Tr}[\rho_{USE} (\Pi_U \otimes \mathbf{1}_{SE})]}$ for some $\Pi > 0$
- ✓ we say that the family \mathcal{S} is **steerable** if there exists a ρ_{USE} such that:
 - 1 for all $\rho_{SE} \in \mathcal{S}$ there exists a $\Pi_U > 0$ such that the above holds, and
 - 2 for all $\Pi_U > 0$, the steered state ρ_{SE}^{Π} is in \mathcal{S}
- ✓ example: if \mathcal{S} is a polytope (e.g., bipartite cq-states) then it's steerable
- ✓ example: the family $\mathcal{S} = \{\rho_S \otimes \bar{\xi}_E\}$ corresponding to the initial factorization condition is steerable
- ✓ **main fact:** bipartite family \mathcal{S} is steerable and Markovian \iff it can be steered from Markovian ρ_{USE} , i.e., $I(U; E|S)_{\rho} = 0$
- ✓ **first corollary:** all previous cases (they all happen to be steerable Markovian families)
- ✓ many other more general constructions are possible too \implies no direct connection between "strength/character of initial correlations" and "existence of CPTP reduced dynamics"
- ✓ **second corollary:** if \mathcal{S} is Markovian, steerable, and such that $\text{Tr}_E[\mathcal{S}]$ contains all states on \mathcal{H}_S , then it must be in factorized form \implies initial factorization condition
- ✓ **extra goody:** we can apply all the tools recently developed for **approximate recoverability** (family is approximately steerable, family is approximately Markov, tripartite state is approximately Markov, etc)

- ✓ data-processing theorem: if there is a process, information always decreases
- ✓ reverse data-processing theorem: if information *always* decreases, then there exists a process
- ✓ data-processing inequality as a “physical principle”: the flow of information determines the evolution
- ✓ analogy with strong (i.e., necessary and sufficient) second law-like statements (e.g., Lieb–Yngvasson formulation of adiabaticity)
- ✓ work in progress: applications to generalized resource theories



è finita la comedia